



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= \frac{8r^2}{\pi^2} \int_0^{\frac{1}{2}\pi} (\frac{1}{8}\pi^2 + \frac{1}{2}\theta^2 - \frac{1}{2}\pi\theta + \frac{1}{2}\cos^2\theta - \frac{1}{2}\pi\sin\theta \cos\theta + \theta\sin\theta \cos\theta) d\theta = \frac{1}{8}\pi r^2.$$

Second Case. The two chords AB and $A'B'$ may be on *different* sides of the diameter XY ; that is, the chord $A'B'$ now becomes the chord MN in position.

\therefore Area $AMNB = U_2 = (\phi + \theta + \sin\phi \cos\phi + \sin\theta \cos\theta)r^2$; and consequently,

$$A_2 = \frac{\int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} U_2 d\theta d\phi d\omega d\psi}{\int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} d\theta d\phi d\omega d\psi}$$

$$= \frac{4r^2}{\pi^2} \int_0^{\frac{1}{2}\pi} (\frac{1}{8}\pi^2 + \frac{1}{2}\pi\theta + \frac{1}{2} + \frac{1}{2}\pi \sin\theta \cos\theta) d\theta = \frac{1}{2}(\pi + 1/\pi)r^2.$$

$\therefore A = \frac{1}{2}(A_1 + A_2) = (\frac{1}{3}\pi + 1/4\pi)r^2$, which is the required average area.

Corollary. If $r=1$, $A = \frac{1}{3}\frac{1}{4}$; that is, the required average is slightly greater than one-third of the given circle.

CALCULUS.

201. Proposed by F. P. MATZ, Sc. D., Ph. D.

$$\text{Solve } \int \int_0^{dy/dx} \frac{dw}{1+w^2} = 0.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

$$\int \int_0^{dy/dx} \frac{dw}{1+w^2} = 0. \quad \int \left[\tan^{-1} w \right]_0^{dy/dx} = 0. \quad \therefore \int \tan^{-1} \frac{dy}{dx} = 0. \quad \therefore \frac{dy}{dx} = 0.$$

$$y = C = \text{a constant.}$$

202. Proposed by W. J. GREENSTREET, M. A., Editor of the Mathematical Gazette, Stroud, England.

Find the complete primitive of $y = 2px + ap^2$. Regard the primitive as the equation giving the arbitrary constant, and if the primitive has equal roots discuss the equation expressing that condition.

Solution by G. W. GREENWOOD, M. A.

Differentiating with regard to x we obtain an equation which may be written

$$p \frac{dx}{dp} + 2x = -2ap.$$

Integrating, we have $3p^2x + 2ap^3 = c$, where c is a constant. Eliminating p between this equation and the original equation, we get

$$a^2c^2 + 2c(2x^3 + 3axy) - y^2(4ay + 3x^2) = 0.$$

This is the primitive equation. If the values of c are equal, then

$$(2x^3 + 3axy)^2 + a^2y^2(4ay + 3x^2) = 0; \text{ i. e., } (x^2 + ay)^3 = 0.$$

When the discriminating equation has a factor cubed, this factor equated to zero gives the cuspidal locus of the given family of curves. In fact, in the present case there is a cusp at the point $(a^{\frac{2}{3}}c^{\frac{1}{3}}, -a^{\frac{1}{3}}c^{\frac{2}{3}})$.

GEOMETRY.

260. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

Perpendiculars to the radius vector are drawn through points on $r = a + b \cos n\theta$. Find the radius of curvature of their envelope at a point at a given distance from the origin.

I. Solution by G. B. M. ZERR, A. M., Ph. D.

$$r = a + b \cos n\theta = R \cos(\phi - \theta) = p \dots \dots (1).$$

$$\text{Differentiating (1) we get, } -b n \sin n\theta = R \sin(\phi - \theta) \dots \dots (2).$$

$$\therefore R^2 = (a + b \cos n\theta)^2 + b^2 n^2 \sin^2 n\theta \dots \dots (3).$$

$$\therefore R^2 = p^2 + b^2 n^2 - n^2(p - a)^2 \dots \dots (4).$$

(3) and (4) are both equations to the envelope.

Radius of curvature $= \rho = R(dR/dp)$.

$$\begin{aligned} \therefore \rho &= p + an^2 - n^2p = r + an^2 - n^2r = an^2 + (1 - n^2)(a + b \cos n\theta) \\ &= a + b \cos n\theta - bn^2 \cos n\theta = a - b(n^2 - 1) \cos n\theta. \end{aligned}$$

II. Solution by G. W. GREENWOOD, M. A.

Let (p, α) be the foot of the perpendicular from the origin upon the tangent at a point P of the envelope. Then at P ,

$$\rho = p + d^2p/da^2 = a + b \cos na - bn^2 \cos na,$$

since

$$p = a + b \cos na.$$

MECHANICS.

182. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

I have a tank, the lower part of which is a hemisphere 22 feet in diameter. The rest is a cylinder 22 feet in diameter, and altitude 28 feet. This tank is connected with the earth by a vertical stand-pipe 10 inches in diameter, 130 feet long, extending 2 feet into the tank. The tank is filled by a $2\frac{1}{2}$ inch pipe 65 feet long, having one right-angled elbow delivering the water into the bottom of the stand-pipe from a steam pump under 96 pounds gauge pressure. How long will it take to fill the pipe?